

References

- Allain, J.Y., 1974. Ecologie des bancs d'Ophiothrix fragilis (Abildgaard)(Echinodermata: Ophiuroidea) dans le golfe normanno-breton. *Cahiers de biologie marine*, 15, pp.255-273.
- Arndt, E.A. 1991. Ecological, physiological and historical aspects of brackish water fauna distribution. In: Estuaries and coasts: Spatial and temporal intercomparisons. Estuarine and Coastal Sciences Association 19 Symposium, ed. by M. Elliott & J.P. Ducrotoy. Olsen and Olsen.
- Bamber, R.N. 1993. Changes in the infauna of a sandy beach. *Journal of Marine Biology and Ecology*, 172: 93-107.
- Basford, D.J., Eleftheriou, A. and Raffaelli, D., 1989. The epifauna of the northern North Sea (56–61 N). *Journal of the Marine Biological Association of the United Kingdom*, 69(2), pp.387-407.
- Basimi, R.A. and Grove, D.J., 1985. Estimates of daily food intake by an inshore population of *Pleuronectes platessa* L. off eastern Anglesey, North Wales. *Journal of Fish Biology*, 27(4), pp.505-520.
- Billett, D.S.M., Bett, B.J., Rice, A.L., Thurston, M.H., Galéron, J., Sibuet, M. and Wolff, G.A., 2001. Long-term change in the megabenthos of the Porcupine Abyssal Plain (NE Atlantic). *Progress in Oceanography*, 50(1-4), pp.325-348.
- Blunden, G., Farnham, W.F., Jephson, N., Fenn, R.H., & Plunket, B.A. 1977. The composition of maerl from the Glenan Islands of South Brittany. *Botanica Marina*, 20: 121-125.
- Blunden, G., Farnham, W.F., Jephson, N., Barwell, C.J., Fenn, R.H., & Plunket, B.A. 1981. The composition of maerl of economic interest in northern Brittany, Cornwall, and Ireland. *Proceedings of the International Seaweed Symposium*, 651-656.
- Bosence, D.W.J. 1979. The factors leading to aggregation and reef formation in *Serpula vermicularis* L. In: Biology and systematics of colonial organisms, ed. by G. Larwood & B.R. Rosen, 11: 299-318. London, Academic Press.
- Brun E (1969) Aggregation of *Ophiothrix fragilis* (Abildgaard) (Echinodermata: Ophiuroidea). *Nytt Mag Zool* 17: 155±160
- Cabioch, L. & Glaçon, R. 1977. Distribution des peuplements benthiques en Manche orientale, du cap d'Antifer à la baie de Somme. C.R. Acad. Sci. Paris, t. 285, sér. D:209-212.
- Dauvin, J.C., Bellan, G., Bellan-Santini, D., Castric, A., Françour, P., Gentil, F., Girard, A., Gofas, S., Mahe, C., Noël, P., & Reviers, B. de. 1994. Typologie des ZNIEFF-MER. Liste des paramètres et des biocoenoses des côtes françaises métropolitaines. 2nd ed. Paris, Muséum National d'Histoire Naturelle, Secrétariat Faune-Flore. (Collection Patrimoines Naturels, Série Patrimoine Ecologique, No. 12.)
- Davies, J. & Hall-Spencer, J. 1996. Mapping of the benthic biotopes in the proposed Sound of Arisaig Special Area of Conservation. Edinburgh, Scottish Natural Heritage. (Scottish Natural Heritage/BioMar Report.)
- Davoult, D. and Gounin, F., 1995. Suspension-feeding activity of a dense *Ophiothrix fragilis* (Abildgaard) population at the water-sediment interface: time coupling of food availability and feeding behaviour of the species. *Estuarine, Coastal and Shelf Science*, 41(5), pp.567-577.
- Degraer, S., Mouton, I., De Neve, L. and Vincx, M., 1999. Community structure and intertidal zonation of the macrobenthos on a macrotidal, ultra-dissipative sandy beach: Summer-winter comparison. *Estuaries*, 22(3), pp.742-752.

- Dewarumez, J.-M., Davoult, D., Anorve, L.E.S & Frontier, S. 1992. Is the "muddy heterogenous sediment assemblage" an ecotone between the pebbles community and the *Abra alba* community in the Southern Bight of the North Sea? *Netherlands Journal of Sea Research*, 30: 229-238.
- Eleftheriou, A. and Basford, D.J., 1989. The macrobenthic infauna of the offshore northern North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 69(1), pp.123-143.
- Ford, E. 1923. Animal communities of the level sea-bottom in the water adjacent to Plymouth. *Journal of the Marine Biological Association of the United Kingdom*, 13: 164-224.
- Freiwald, A., Fossa, J.H., Grehan, A., Koslow, T. and Roberts, J.M., 2004. Cold-water coral reefs: out of sight-no longer out of mind. UNEP-WCMC.
- Gage, J.D., 1986. The benthic fauna of the Rockall Trough: regional distribution and bathymetric zonation. *Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences*, 88, pp.159-174.
- Gameson, A.L.H. ed., 1982. The quality of the Humber estuary, 1961-1981. Yorkshire Water Authority.
- Glémarec, M. 1973. The benthic communities of the European North Atlantic continental shelf. *Oceanography and Marine Biology. An Annual Review*, 11: 263-289.
- Hiscock, K. 1984. Sublittoral survey of Bardsey and the Lleyn peninsula. August 13th to 27th, 1983. (Contractor: Field Studies Council, Oil Pollution Research Unit, Pembroke.) Nature Conservancy Council, CSD Report, No. 612.
- Hughes, D.J. and Gage, J.D., 2004. Benthic metazoan biomass, community structure and bioturbation at three contrasting deep-water sites on the northwest European continental margin. *Progress in Oceanography*, 63(1-2), pp.29-55.
- Husebø, Å., Nøttestad, L., Fosså, J.H., Furevik, D.M. and Jørgensen, S.B., 2002. Distribution and abundance of fish in deep-sea coral habitats. *Hydrobiologia*, 471(1), pp.91-99.
- Irvine, L.M., & Chamberlain, Y.M. 1994. Seaweeds of the British Isles. Volume 1 Rhodophyta. Part2B Corallinales, Hildenbrandiales. London, HMSO for The Natural History Museum.
- Jones, N.S. 1950. Marine bottom communities. *Biological Reviews*, 25: 283-313.
- Jones, N.S. 1951. The bottom fauna off the south of the Isle of Man. *Journal of Animal Ecology*, 20: 132-144.
- Kröncke, I., 1990. Macrofauna standing stock of the Dogger Bank. A comparison: II. 1951–1952 versus 1985–1987 are changes in the community of the northeastern part of the Dogger Bank due to environmental changes? *Netherlands Journal of Sea Research*, 25(1-2), pp.189-198.
- Kühne, S. & Rachnor, E. 1996. The macrofauna of a stony sand area in the German Bight (North Sea). *Helgoländer Meeresuntersuchungen*. 50: 433-452.
- Künitzer, A., Basford, D., Craeymeersch, J.A., Dewarumez, J.M., Derjes, J., Duineveld, G.C.A., Eleftheriou, A., Heip, C., Herman, P., Kingston, P., Neirmann, U., Rachor, E., Rumohr, H., & Wilde, P.A.J.de. 1992. The benthic infauna of the North Sea: species distribution and assemblages. *ICES Journal of Marine Science*, 49: 127-143.
- Lavaleye, M.S.S., Duineveld, G.C.A., Berghuis, E.M., Kok, A. and Witbaard, R., 2002. A comparison between the megafauna communities on the NW Iberian and Celtic continental margins—effects of coastal upwelling. *Progress in Oceanography*, 52(2-4), pp.459-476.

- Lockwood, S.J., 1980. Density-dependent mortality in 0-group plaice (*Pleuronectes platessa* L.) populations. *ICES Journal of Marine Science*, 39(2), pp.148-153.
- Long, D. 2006. BGS detailed explanation of seabed sediment modified folk classification. Available at http://www.searchmesh.net/PDF/GMHM3_Detailed_explanation_of_seabed_sediment_classification.pdf
- Macer, C.T., 1967. The food web in Red Wharf Bay (North Wales) with particular reference to young plaice (*Pleuronectes platessa*). *Helgoländer Wissenschaftliche Meeresuntersuchungen*, 15(1), pp.560-573.
- Mackie, A.S.Y. 1990. Offshore benthic communities of the Irish Sea. In: *The Irish Sea: an environmental review. Part 1: nature conservation*, ed. by Irish Sea Study Group, 169-218. Liverpool, Liverpool University Press for Irish Sea Study Group.
- Mackie, A.S.Y., Oliver, P.G., & Rees, E.I.S. 1995. Benthic biodiversity in the southern Irish sea. Unpublished, *Studies in Marine Biodiversity and Systematics from the National Museum of Wales*. (Biomor report, No. 1.)
- McIntyre, A.D. 1961. Quantitative differences in the fauna of boreal mud associations. *Journal of the Marine Biological Association of the United Kingdom*, 41: 599-616.
- Moore, C.G. 1996. The distribution of serpulid reefs in Loch Creran, Argyll. *Scottish Natural Heritage Research, Survey and Monitoring Report*, No. 53.
- Mortensen, P.B., Hovland, M., Brattegard, T. and Farestveit, R., 1995. Deep water bioherms of the scleractinian coral *Lophelia pertusa* (L.) at 64 N on the Norwegian shelf: structure and associated megafauna. *Sarsia*, 80(2), pp.145-158.
- Pearson, T.H., & Rosenberg, R. 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanography and Marine Biology. An Annual Review*, 16: 229-311.
- Pearson, T.H., Mannvik, Hans-Petter, Evans, R., & Falk-Petersen, Falk. 1996. The benthic communities of the Snorre Field in the Northern North Sea. 1. The distribution and structure of communities in undisturbed sediments. *Journal of Sea Research* 35(4): 301-314.
- Petersen, C.G.J. 1918. The sea bottom and its production of fish-food. A survey of the work done in connection with valuation of the Denmark waters from 1883-1917. *Report of the Danish Biological Station*, 25: 1-62.
- Picton, B.E. 1993. A field guide to the shallow-water echinoderms of the British Isles. London, Immel Publishing for Marine Conservation Society.
- Picton, B.E., Emblow, C.S., Morrow, C.C., Sides, E.M., & Costello, M.J. 1994. Marine communities of the Mulroy Bay and Lough Swilly area, north-west Ireland, with an assessment of their nature conservation importance. Unpublished, Environmental Sciences Unit, Trinity College. (Field Survey Report.)
- Rodwell, J.S. ed. 1995. *British plant communities. Volume 4. Aquatic communities, swamps and tall herb fens*. Cambridge, Cambridge University Press.
- Rodwell, J.S. 2000. *British plant communities. Volume 5: Maritime communities and vegetation of open habitats*. Cambridge, Cambridge University Press.
- Salzwedel, H., Rachor, E., & Gerdes, D. 1985. Benthic macrofauna communities in the German Bight. *Veröffentlichungen des Institut für Meeresforschung in Bremerhaven*, 20: 199-267.
- Stephen, A.C. 1923. Preliminary survey of the Scottish waters of the North Sea by the Petersen grab. *Scientific Investigations of the Fisheries Division of the Scottish Home Department*, 11.

- Tait, R.V. & Dipper, R.A., 1998. *Elements of Marine Ecology*. Reed Elsevier
- Thorson, G. 1934. Contributions to the animal ecology of the Scoresby Sound Fjord complex (east Greenland). *Meddelelser om Grønland. Kommissionen for Videnskabelige Undersøgelser I Grønland*, 100 (3): 1-67.
- Thorson, G. 1957. Bottom communities (sublittoral or shallow shelf). *Memoirs of the Geological Society of America*, 67: 461-534.
- Tittley, I., Farnham, W.F., Fletcher, R.L., Morell, S. and Bishop, G., 1985. Sublittoral seaweed assemblages of some northern Atlantic islands. *Progress in Underwater Science*, 10, pp.39-52.
- Tittley, I., Gilliland, P. and Pound, D., 1998. Marine flora of the Thanet Coast cSAC: a conservation and management perspective. Changes in the marine flora of the North Sea. Centre for Research into Coastal Issues, Scarborough, pp.65-75.
- Tittley, I., Spurrier, C.J.H. and Chimonides, P.J., 2004. Thanet Intertidal Survey: Assessment of Favourable Condition of Reef and Sea-cave Features in the Thanet Coast CSAC: January 2002. English Nature.
- Vanosmael, C., Willems, K.A., Claeys, D., Vincx, M. and Heip, C., 1982. Macrobenthos of a sublittoral sandbank in the Southern Bight of the North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 62(3), pp.521-534.
- Vorberg, R., 2000. Effects of shrimp fisheries on reefs of *Sabellaria spinulosa* (Polychaeta). *ICES Journal of Marine Science*, 57(5), pp.1416-1420.
- Warner, G.F., 1971. On the ecology of a dense bed of the brittle-star *Ophiotrix fragilis*. *Journal of the Marine Biological Association of the United Kingdom*, 51(2), pp.267-282.
- Warren, P.J. 1977. The Nephrops fishery off north-east England. *Annales Biologiques*, 34: 234-235
- Warwick, R.M., & Davies, J.R. 1977. The distribution of sublittoral macrofauna communities in the Bristol Channel in relation to the substrate. *Estuarine and Coastal Marine Science*, 5: 267-288.
- Warwick, R.M., & Uncles, R.J. 1980. Distribution of benthic macrofauna associations in the Bristol Channel in relation to tidal stress. *Marine Ecology (Progress Series)*, 3: 97-103.
- Willems, K.A., Vanosmael, C., Claeys, D., Vincx, M. and Heip, C., 1982. Benthos of a sublittoral sandbank in the southern bight of the North Sea: general considerations. *Journal of the Marine Biological Association of the United Kingdom*, 62(3), pp.549-557.
- Wilson, D.P., 1929. The larvae of the British sabellarians. *Journal of the Marine Biological Association of the United Kingdom*, 16(1), pp.221-268.
- Wilson, J. G. 1979. The distribution of the coral *Lophelia pertusa* (L.) (*L. prolifera* (Pallas)) in the north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom* 59, 149–164.
- Wolff, W.J. 1973. The estuary as a habitat. An analysis of data on the soft-bottom macrofauna on the estuarine area of the rivers Rhine, Meuse and Scheldt. Zoologischer Verhandelingen, (Delta Institute for Hydrobiological Research, Communication, No. 106): 126